



Travis G. Maak, M.D.
 590 Wakara Way
 Salt Lake City, UT 84108
 Tel: (801) 587-7109
 Fax: (801)587-7112
 Lic. # 8234797-1205

Physical Therapy Prescription Osteochondral Allograft

Patient Name: _____ **Today's Date:** _____ **Surgery Date:** _____

Dx: s/p (LEFT / RIGHT) KNEE Osteochondral Allograft (MFC, LFC)

Modalities:

Weeks 0-4

- ___ Strict TTWB x 4 weeks with brace locked at 0 degrees, may progress to FWB by 6 weeks
- ___ Hinged, double-upright brace in full extension
- ___ Quadriceps and Adductor isometrics
- ___ SLR's into flexion, extension, abduction and adduction (AAROM→AROM)
- ___ Passive (CPM) and AAROM (ROM limited to 0-90° knee flexion for 4 weeks)
- ___ Manual and self-mobilizations for Hamstrings, Gastrocnemius, Hip flexors and ITB/lateral retinaculum
- ___ Gentle Patella, fibular head and scar mobilization
- ___ NMES for Quadriceps re-education daily
- ___ Cryotherapy and cryokinetics for pain

Weeks 4-6

- ___ Continue hinged, double-upright brace in full extension
- ___ Progressive partial WB to full WB by week 6, brace still must be locked at 0 degrees
- ___ Initiate PWB closed kinetic chain strengthening, short arc
- ___ Continue core strength and stability
- ___ Begin gentle Quadriceps stretching as tolerated
 (Concentrating on proximal attachment – limiting knee flexion to less than 120°)
- ___ Continue gentle Patella, fibular head and scar mobilization
- ___ Begin proprioception exercise on stable surface

Weeks 6-12

- ___ Discontinue Hinged, double-upright brace in full extension
- ___ Full weight bearing
- ___ Begin gait training and standing proprioceptive training on unstable surface
- ___ Progress closed kinetic chain strengthening for lower extremity
- ___ Progress OKC and CKC strengthening for hip
- ___ Continue gentle Patella, fibular head and scar mobilization
- ___ Continue core and hip strength and endurance

Weeks 12-24

- ___ Continue gait training and proprioceptive training
- ___ Progress closed and open kinetic chain strengthening for lower extremity
 (Avoid hyperflexion at knee and full open kinetic chain knee extension)
- ___ Continue lower extremity strengthening
- ___ Begin light jogging, progress to light recreational activities by 20-24 weeks
- ___ Continue NMES for Quadriceps re-education PRN
- ___ Continue Cryotherapy for pain/edema/effusion PRN
- ___ Begin agility drills in single plane, frontal and sagittal
- ___ Pivoting to begin at 4.5 months
- ___ Assess posture and functional movement patterns. Corrective exercise as needed.

Frequency & Duration: (circle one) 1-2 2-3 x/week for _____ weeks Home Program

**Please send progress notes.

Physician's Signature: _____ **M.D.**